

Spaces with Ricci curvature bounded from below

Nicola Gigli

February 24, 2015

Topics

- 1) On the definition of spaces with Ricci curvature bounded from below
- 2) Analytic properties of $RCD^*(K, N)$ spaces
- 3) Geometric properties of $RCD^*(K, N)$ spaces
- 4 More on the differential structure of metric measure spaces

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

The Abresch-Gromoll inequality

On a Riemannian manifold with $\text{Ric} \geq K$ and $\dim \leq N$ we have

$$E(x) \leq f_{K,N}(h(x)), \quad \text{provided} \quad h(x) \leq \frac{\min\{d(x, \gamma_0), d(x, \gamma_1)\}}{2}$$

for some (explicitly given) $f_{K,N}$ satisfying

$$\lim_{h \downarrow 0} \frac{f_{K,N}(h)}{h} = 0.$$

Ingredients of the proof

Laplacian comparison estimates for the distance

Linearity of the Laplacian

Weak maximum principle

The non-smooth case

Repeating verbatim the proof on $RCD^*(K, N)$ spaces we obtain:

Thm. (G., Mosconi '12) The Abresch-Gromoll inequality holds in the non-smooth setting in the same form as in the smooth one.

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

The splitting theorem

Thm. (Cheeger-Gromoll '71)

Let M be a Riemannian manifold with $\text{Ric} \geq 0$ which contains a line.
Then $M = N \times \mathbb{R}$ for some Riemannian manifold N .

The almost splitting

Thm. (Cheeger-Colding '96) Let M be a Riemannian manifold with $\text{Ric} \geq -\varepsilon$ which contains a geodesic with length L , with $\varepsilon, L^{-1} \ll 1$

Then 'a big portion of M is mGH-close to a product'

The almost splitting

Thm. (Cheeger-Colding '96) Let M be a Riemannian manifold with $\text{Ric} \geq -\varepsilon$ which contains a geodesic with length L , with $\varepsilon, L^{-1} \ll 1$

Then 'a big portion of M is mGH-close to a product'

Cor. (Splitting for limit spaces) Let (X, d, \mathfrak{m}) be a pmGH of Riemannian manifolds (M_n) with uniformly bounded dimension and with $\text{Ric}(M_n) \geq -\varepsilon_n$, where $\varepsilon_n \downarrow 0$.

Assume that X contains a line. Then it splits off a factor \mathbb{R}

The non-smooth splitting

Thm. (G. '13) Let (X, d, \mathbf{m}) be an $RCD^*(0, N)$ space containing a line. Then there is a space (X', d', \mathbf{m}') such that

$$(X, d, \mathbf{m}) \text{ is isomorphic to } (X' \times \mathbb{R}, d' \otimes d_{\text{Eucl}}, \mathbf{m}' \times \mathcal{L}^1)$$

where

$$(d' \otimes d_{\text{Eucl}})((x', t), (y', s)) := \sqrt{d'(x', y')^2 + |t - s|^2}$$

Moreover:

- ▶ If $N \geq 2$ then (X', d', \mathbf{m}') is an $RCD^*(0, N - 1)$ space
- ▶ If $N \in [1, 2)$ then X' contains only one point

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

The Busemann function

Let $\gamma : [0, \infty) \rightarrow M$ an half line.

The Busemann function b associated to it is

$$b(x) := \lim_{t \rightarrow +\infty} t - d(x, \gamma_t) = \sup_{t \geq 0} t - d(x, \gamma_t)$$

The Busemann function

Let $\gamma : [0, \infty) \rightarrow M$ an half line.

The Busemann function b associated to it is

$$b(x) := \lim_{t \rightarrow +\infty} t - d(x, \gamma_t) = \sup_{t \geq 0} t - d(x, \gamma_t)$$

If $\gamma : (-\infty, +\infty) \rightarrow M$ is a line we can associate to it 2 Busemann functions

$$b^+(x) := \lim_{t \rightarrow +\infty} t - d(x, \gamma_t)$$

$$b^-(x) := \lim_{t \rightarrow +\infty} t - d(x, \gamma_{-t})$$

Effect of $\text{Ric} \geq 0$ on the Busemann function for an half line

If $\text{Ric} \geq 0$ and $\bar{x} \in M$

$$\Delta \frac{d^2(\cdot, \bar{x})}{2} \leq \dim(M)$$

Hence

$$\Delta d(\cdot, \gamma_t) \leq \frac{\dim(M)}{d(\cdot, \gamma_t)}$$

Passing to the limit we obtain

$$\Delta b \geq 0,$$

i.e. the b is subharmonic.

What for the Busemann function for a line

b^+ and b^- are subharmonic, thus so is $b^+ + b^-$.
The triangle inequality gives

$$b^+ + b^- \leq 0$$

and the fact that γ is a line ensures that

$$(b^+ + b^-)(\gamma_0) = 0$$

hence (strong maximum principle) it holds

$$b^+ + b^- \equiv 0$$

and in particular b^+ and b^- are harmonic

Use of the Bochner equality and inequality

For any f smooth it holds

$$\begin{aligned}\Delta \frac{|\nabla f|^2}{2} &= \|\text{Hess } f\|_{\text{HS}}^2 + \nabla f \cdot \nabla \Delta f + \text{Ric}(\nabla f, \nabla f) \\ &\geq \frac{(\Delta f)^2}{\dim(M)} + \nabla f \cdot \nabla \Delta f\end{aligned}$$

Use of the Bochner equality and inequality

For any f smooth it holds

$$\begin{aligned}\Delta \frac{|\nabla f|^2}{2} &= \|\text{Hess } f\|_{\text{HS}}^2 + \nabla f \cdot \nabla \Delta f + \text{Ric}(\nabla f, \nabla f) \\ &\geq \frac{(\Delta f)^2}{\dim(M)} + \nabla f \cdot \nabla \Delta f\end{aligned}$$

For b^+ we have $|\nabla b^+| \equiv 1$ and $\Delta b^+ \equiv 0$ and thus the equality

$$\Delta \frac{|\nabla b^+|^2}{2} = \frac{(\Delta b^+)^2}{\dim(M)} + \nabla b^+ \cdot \nabla \Delta b^+$$

Use of the Bochner equality and inequality

For any f smooth it holds

$$\begin{aligned}\Delta \frac{|\nabla f|^2}{2} &= \|\text{Hess } f\|_{\text{HS}}^2 + \nabla f \cdot \nabla \Delta f + \text{Ric}(\nabla f, \nabla f) \\ &\geq \frac{(\Delta f)^2}{\dim(M)} + \nabla f \cdot \nabla \Delta f\end{aligned}$$

For b^+ we have $|\nabla b^+| \equiv 1$ and $\Delta b^+ \equiv 0$ and thus the equality

$$\Delta \frac{|\nabla b^+|^2}{2} = \frac{(\Delta b^+)^2}{\dim(M)} + \nabla b^+ \cdot \nabla \Delta b^+$$

which yields

$$\|\text{Hess } b^+\|_{\text{HS}}^2 \equiv \frac{(\Delta b^+)^2}{\dim(M)} \equiv 0$$

i.e. b^+ is both convex and concave.

Isometries via gradient flows

Since b^+ is convex, its gradient flow contracts distances.

Since $b^+ = -b^-$ is concave, its gradient flow expands distances.

Thus the gradient flow of b^+ produces a 1-parameter family of isometries.

Conclusion of the argument

Put $N := \{b^+ = 0\}$

Conclusion of the argument

Put $N := \{b^+ = 0\}$

For $x \in N$ and $v \in T_x N$ it is obvious that $v \cdot \nabla b^+(x) = 0$ and the conclusion follows from the fact the gradient flow of b^+ is a 1-parameter family of isometries.

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

Harmonicity of b^\pm

As in the smooth case, from the Laplacian comparison estimate we deduce

$$\Delta b^\pm \geq 0,$$

and using the strong maximum principle (Bjorn-Bjorn '07) we obtain

$$b^+ + b^- = 0$$

i.e.

$$\Delta b^\pm = 0$$

Gradient flow of b^\pm and geodesics

For every $t \in \mathbb{R}$ the function tb^+ is c -concave and

$$(tb^+)^c = tb^- - \frac{t^2}{2}$$

Gradient flow of b^\pm and geodesics

For every $t \in \mathbb{R}$ the function tb^+ is c -concave and

$$(tb^+)^c = tb^- - \frac{t^2}{2}$$

Hence from existence and uniqueness of optimal maps we deduce that

for \mathbf{m} -a.e. $x \in X$ there is a unique $F_t(x) \in \partial^c(tb^+)(x)$

Gradient flow of b^\pm and geodesics

For every $t \in \mathbb{R}$ the function tb^+ is c -concave and

$$(tb^+)^c = tb^- - \frac{t^2}{2}$$

Hence from existence and uniqueness of optimal maps we deduce that

for \mathbf{m} -a.e. $x \in X$ there is a unique $F_t(x) \in \partial^c(tb^+)(x)$

and we can check that for \mathbf{m} -a.e. x we have

$$\begin{aligned} t &\mapsto F_t(x), && \text{is a line} \\ F_{t+s}(x) &= F_t(F_s(x)), && \forall t, s \in \mathbb{R} \end{aligned}$$

Measure preservation

For every $\mu = \rho \mathbf{m} \ll \mathbf{m}$ the map $[0, 1] \ni t \mapsto (F_t)_\# \mu$ is a W_2 -geodesic induced by \mathbf{b}^+ .

Arguing as in the proof of the Laplacian comparison estimates we deduce

$$\frac{1}{t} (\mathcal{U}_N((F_t)_\# \mu) - \mathcal{U}_N(\mu)) \geq -\frac{1}{N} \int \nabla(\rho^{1-\frac{1}{N}}) \cdot \nabla \mathbf{b}^+ \, d\mathbf{m} = 0$$

Measure preservation

For every $\mu = \rho \mathbf{m} \ll \mathbf{m}$ the map $[0, 1] \ni t \mapsto (F_t)_\# \mu$ is a W_2 -geodesic induced by b^+ .

Arguing as in the proof of the Laplacian comparison estimates we deduce

$$\frac{1}{t} (\mathcal{U}_N((F_t)_\# \mu) - \mathcal{U}_N(\mu)) \geq -\frac{1}{N} \int \nabla(\rho^{1-\frac{1}{N}}) \cdot \nabla b^+ \, d\mathbf{m} = 0$$

Switching b^+ and b^- we deduce

$$\mathcal{U}_N((F_t)_\# \mu) = \mathcal{U}_N(\mu), \quad \forall t \in \mathbb{R}$$

and thus

$$(F_t)_\# \mathbf{m} = \mathbf{m}, \quad \forall t \in \mathbb{R}$$

How to use Bochner inequality

We have

$$\Delta \frac{|\nabla f|^2}{2} \geq \nabla f \cdot \nabla \Delta f$$

How to use Bochner inequality

We have

$$\Delta \frac{|\nabla f|^2}{2} \geq \nabla f \cdot \nabla \Delta f$$

Write it with $b^+ + \varepsilon f$ in place of f we obtain

$$\Delta(\nabla b^+ \cdot \nabla f) = \nabla b^+ \cdot \nabla \Delta f$$

for every f 'smooth enough'.

How to use Bochner inequality

We have

$$\Delta \frac{|\nabla f|^2}{2} \geq \nabla f \cdot \nabla \Delta f$$

Write it with $b^+ + \varepsilon f$ in place of f we obtain

$$\Delta(\nabla b^+ \cdot \nabla f) = \nabla b^+ \cdot \nabla \Delta f$$

for every f 'smooth enough'.

Using this identity in computing $\frac{d}{dt} \frac{1}{2} \int |D(f \circ F_t)|^2 d\mathbf{m}$ we get

$$\frac{d}{dt} \frac{1}{2} \int |D(f \circ F_t)|^2 d\mathbf{m} = 0$$

and thus

$$t \mapsto \frac{1}{2} \int |D(f \circ F_t)|^2 d\mathbf{m} \quad \text{is constant}$$

Isomorphisms by duality

Let (X_1, d_1) , (X_2, d_2) be metric spaces and $T : X_1 \rightarrow X_2$ invertible.

Isomorphisms by duality

Let (X_1, d_1) , (X_2, d_2) be metric spaces and $T : X_1 \rightarrow X_2$ invertible.
Then T is an isometry if and only if

$$\text{Lip}_{X_1}(f \circ T) = \text{Lip}_{X_2}(f), \quad \forall f : X_2 \rightarrow \mathbb{R}$$

Isomorphisms by duality

Let (X_1, d_1) , (X_2, d_2) be metric spaces and $T : X_1 \rightarrow X_2$ invertible.
Then T is an isometry if and only if

$$\text{Lip}_{X_1}(f \circ T) = \text{Lip}_{X_2}(f), \quad \forall f : X_2 \rightarrow \mathbb{R}$$

Let (X_1, d_1, \mathbf{m}_1) , (X_2, d_2, \mathbf{m}_2) be $RCD^*(K, \infty)$ spaces and $T : X_1 \rightarrow X_2$ Borel and a.e. invertible.

Isomorphisms by duality

Let (X_1, d_1) , (X_2, d_2) be metric spaces and $T : X_1 \rightarrow X_2$ invertible.
Then T is an isometry if and only if

$$\text{Lip}_{X_1}(f \circ T) = \text{Lip}_{X_2}(f), \quad \forall f : X_2 \rightarrow \mathbb{R}$$

Let (X_1, d_1, \mathbf{m}_1) , (X_2, d_2, \mathbf{m}_2) be $RCD^*(K, \infty)$ spaces and $T : X_1 \rightarrow X_2$ Borel and a.e. invertible.

Then T is (up to a modification on a negligible set) an isomorphism if and only if

$$\|f \circ T\|_{W^{1,2}(X_1)} = \|f\|_{W^{1,2}(X_2)}, \quad \forall f : X_2 \rightarrow \mathbb{R}$$

The quotient space

We deduce that the F_t 's have representatives which are isometries.

The quotient space

We deduce that the F_t 's have representatives which are isometries.

We can then declare $x \sim y$ if $x = F_t(y)$ for some $t \in \mathbb{R}$, put $X' := X / \sim$ and define

$$d'(\pi(x), \pi(y)) := \inf_{t \in \mathbb{R}} d(x, F_t(y)) \quad \forall x, y \in X$$

and

$$\mathbf{m}'(E) := \mathbf{m}(\pi^{-1}(E)) \cap \mathbf{b}^{-1}([0, 1]) \quad \forall E \subset X' \text{ Borel}$$

The quotient space

We deduce that the F_t 's have representatives which are isometries.

We can then declare $x \sim y$ if $x = F_t(y)$ for some $t \in \mathbb{R}$, put $X' := X / \sim$ and define

$$d'(\pi(x), \pi(y)) := \inf_{t \in \mathbb{R}} d(x, F_t(y)) \quad \forall x, y \in X$$

and

$$\mathbf{m}'(E) := \mathbf{m}(\pi^{-1}(E)) \cap \mathbf{b}^{-1}([0, 1]) \quad \forall E \subset X' \text{ Borel}$$

Define $\iota : X' \rightarrow X$ as

$$\iota(x') = x \quad \text{if} \quad \pi(x) = x' \quad \text{and} \quad \mathbf{b}^+(x) = 0.$$

The quotient space

We deduce that the F_t 's have representatives which are isometries.

We can then declare $x \sim y$ if $x = F_t(y)$ for some $t \in \mathbb{R}$, put $X' := X / \sim$ and define

$$d'(\pi(x), \pi(y)) := \inf_{t \in \mathbb{R}} d(x, F_t(y)) \quad \forall x, y \in X$$

and

$$\mathbf{m}'(E) := \mathbf{m}(\pi^{-1}(E)) \cap \mathbf{b}^{-1}([0, 1]) \quad \forall E \subset X' \text{ Borel}$$

Define $\iota : X' \rightarrow X$ as

$$\iota(x') = x \quad \text{if} \quad \pi(x) = x' \quad \text{and} \quad \mathbf{b}^+(x) = 0.$$

Problem: is ι an isometry?

How to gain C^1 regularity

Let (μ_t) be a geodesic such that $\mu_t \leq C\mathbf{m}$ for every $t \in [0, 1]$ and φ_t
s.t. $-(1-t)\varphi_t$ is a Kantorovich potential from μ_t to μ_1

How to gain C^1 regularity

Let (μ_t) be a geodesic such that $\mu_t \leq C\mathbf{m}$ for every $t \in [0, 1]$ and φ_t s.t. $-(1-t)\varphi_t$ is a Kantorovich potential from μ_t to μ_1

Then:

For $f \in W^{1,2}(X)$ the map $t \mapsto \int f d\mu_t$ is C^1 and

$$\frac{d}{dt} \int f d\mu_t = \int \nabla f \cdot \nabla \varphi_t d\mu_t, \quad \forall t \in [0, 1]$$

How to gain C^1 regularity

Let (μ_t) be a geodesic such that $\mu_t \leq C\mathbf{m}$ for every $t \in [0, 1]$ and φ_t s.t. $-(1-t)\varphi_t$ is a Kantorovich potential from μ_t to μ_1

Then:

For $f \in W^{1,2}(X)$ the map $t \mapsto \int f d\mu_t$ is C^1 and

$$\frac{d}{dt} \int f d\mu_t = \int \nabla f \cdot \nabla \varphi_t d\mu_t, \quad \forall t \in [0, 1]$$

For $\nu \in \mathcal{P}_2(X)$ the map $t \mapsto \frac{1}{2} W_2^2(\mu_t, \nu)$ is C^1 and

$$\frac{d}{dt} \frac{1}{2} W_2^2(\mu_t, \nu) = \int \nabla \phi_t \cdot \nabla \varphi_t d\mu_t, \quad \forall t \in [0, 1]$$

where ϕ_t is a Kantorovich potential from μ_t to ν .

Basic properties of (X', d', \mathbf{m}')

Arguing as in the smooth case but at the level of probability measures $\mu, \nu \leq \mathbf{Cm}$ we deduce that

the minimum of $t \mapsto \frac{1}{2} W_2^2((F_t)_\# \mu, \nu)$

is attained at that t_0 such that $\int b^+ d(F_{t_0})_\# \mu = \int b^+ d\nu$

Basic properties of (X', d', \mathbf{m}')

Arguing as in the smooth case but at the level of probability measures $\mu, \nu \leq \mathbf{Cm}$ we deduce that

the minimum of $t \mapsto \frac{1}{2} W_2^2((F_t)_\# \mu, \nu)$

is attained at that t_0 such that $\int b^+ d(F_{t_0})_\# \mu = \int b^+ d\nu$

Picking μ, ν with support going to a point we conclude that ι is indeed an isometric embedding.

Basic properties of (X', d', \mathbf{m}')

Arguing as in the smooth case but at the level of probability measures $\mu, \nu \leq C\mathbf{m}$ we deduce that

the minimum of $t \mapsto \frac{1}{2} W_2^2((F_t)_\# \mu, \nu)$

is attained at that t_0 such that $\int b^+ d(F_{t_0})_\# \mu = \int b^+ d\nu$

Picking μ, ν with support going to a point we conclude that ι is indeed an isometric embedding.

It is then easy to see that (X', d', \mathbf{m}') is an $RCD^*(0, N)$ space

What remains to show

- (1) That X is isometric to $X' \times \mathbb{R}$
- (2) That X' is an $RCD^*(0, N - 1)$ space

What remains to show

- (1) That X is isometric to $X' \times \mathbb{R}$
- (2) That X' is an $RCD^*(0, N - 1)$ space

The first follows using again the duality with Sobolev functions

The second by a general dimension-reduction argument introduced by [Cavalletti-Sturm](#)

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

The statement

Thm. (Ketterer '13) Let (X, d_X, \mathbf{m}_X) be a metric measure space and (Y, d_Y, \mathbf{m}_Y) be the N -cone built over it. Then TFAE:

- ▶ (X, d_X, \mathbf{m}_X) is an $RCD^*(N-1, N)$ space
- ▶ (Y, d_Y, \mathbf{m}_Y) is an $RCD^*(0, N+1)$ space

Proved via the study of the Bochner inequality

The statement

Thm. (Ketterer '13) Let (X, d_X, \mathbf{m}_X) be a metric measure space and (Y, d_Y, \mathbf{m}_Y) be the N -cone built over it. Then TFAE:

- ▶ (X, d_X, \mathbf{m}_X) is an $RCD^*(N-1, N)$ space
- ▶ (Y, d_Y, \mathbf{m}_Y) is an $RCD^*(0, N+1)$ space

Proved via the study of the Bochner inequality

Corollary - maximal diameter theorem (Ketterer '13) Let (X, d, \mathbf{m}) be a $RCD^*(N-1, N)$ space having two points at distance π . Then it is isomorphic to a spherical suspension over a space (X', d', \mathbf{m}') . Moreover:

- ▶ If $N \geq 2$ then (X', d', \mathbf{m}') is an $RCD^*(N-2, N-1)$ space
- ▶ If $N \in [1, 2)$ then X' contains either only one point or two points at distance π .

Geometric properties of $RCD^*(K, N)$ spaces

- ▶ The Abresch-Gromoll inequality
- ▶ The splitting theorem
 - ▶ Statement
 - ▶ The proof in the smooth case
 - ▶ The proof in the non-smooth case
- ▶ Cones and lower Ricci bounds
- ▶ Rectifiability results

About rectifiability

Theorem (Mondino-Naber '14 / in progress) Let (X, d, \mathbf{m}) be a $\text{RCD}^*(K, N)$ space and $\varepsilon > 0$.

Then there is a Borel partition (A_n) of X and maps $\varphi_n : A_n \rightarrow \mathbb{R}^{d_n}$ with $d_n \leq N$ such that

$$\text{Lip}(\varphi_n), \text{Lip}(\varphi_n^{-1}) \leq 1 + \varepsilon$$

and putting $\mu_n := (\varphi_n)_*(\mathbf{m}|_{A_n})$ we have $\mu_n = \rho_n \mathcal{L}^{d_n}$ with

$$\text{ess-sup}_{\varphi_n(A_n)} \rho_n - \text{ess-inf}_{\varphi_n(A_n)} \rho_n \leq \varepsilon$$

Thank you