

# Li-Yau type estimates and Harnack inequalities by Stochastic Analysis

Anton Thalmaier  
University of Luxembourg

HIM Bonn  
October 10, 2007

### *Heat equation on a manifold*

- Let  $u$  be a (positive) solution of the heat equation on a Riemannian manifold:

$$\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$$

- How to estimate the gradient  $|\nabla u|(x)$ ?

- **Li-Yau** Let  $M$  be complete and  $\text{Ric} \geq -K$ ,  $K \geq 0$ .  
 Let  $u$  be a strictly positive solution of  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  on  $M \times \mathbb{R}_+$   
 and let  $a > 1$ . Then

$$\left( \frac{|\nabla u|}{u} \right)^2 (x, t) - a \frac{\partial_t u}{u}(x, t) \leq c(n, a) \left[ K + \frac{1}{t} \right]$$

[ $a=1$  is possible if  $\text{Ric} \geq 0$ , i.e.  $K = 0$ ].

- **Hamilton** Let  $M$  be complete and  $\text{Ric} \geq -K$ ,  $K \geq 0$ .  
 Let  $u$  be a strictly positive solution of  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  on  $M \times \mathbb{R}_+$   
 and suppose  $u \leq M$  where  $M$  is a real constant. Then

$$\left( \frac{|\nabla u|}{u} \right)^2 (x, t) \leq 2 \left[ K + \frac{1}{t} \right] \log \frac{M}{u(x, t)}.$$

- **Li-Yau** (*Localized version*)

Let  $u > 0$  solve  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  on  $B(x, R) \times [0, t]$ . Then

$$\left( \frac{|\nabla u|}{u} \right)^2 (x, t) - a \frac{\partial_t u}{u} (x, t) \leq c(n, a) \left[ K + \frac{1}{t} + \frac{1}{R^2} \right].$$

- **Hamilton** (*Localized version*)

Let  $0 < u \leq M$  solve  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  on  $B(x, R) \times [0, t]$ . Then

$$\frac{|\nabla u|}{u} (x, t) \leq c(n) \left[ \sqrt{K} + \frac{1}{\sqrt{t}} + \frac{1}{R} \right] \left( 1 + \log \frac{M}{u(x, t)} \right).$$

## Harmonic functions

- Let  $u$  be a (positive) harmonic function on some domain in a Riemannian manifold:

$$\Delta u = 0.$$

- Cheng-Yau**

Let  $M$  be complete and  $D \subset M$  be open, relatively compact. Let  $u : D \rightarrow \mathbb{R}$  be harmonic and strictly positive. Then

$$\frac{|\nabla u|}{u}(x) \leq c(n) \left[ \sqrt{K} + \frac{1}{r(x)} \right]$$

if  $\text{Ric}|_D \geq -K$ ,  $K \geq 0$ .

## *Heat equation with respect moving Riemannian metrics*

- **Perelman** heat equation under backward/forward Ricci flow
- Let  $u$  be a (positive) solution of the following heat equation:

$$\begin{cases} \frac{\partial}{\partial t} u = \frac{1}{2} \Delta_{g(t)} u \\ \frac{\partial}{\partial t} g_t = \pm \text{Ric}_{g(t)} \end{cases}$$

- Find estimates for  $\frac{|\nabla u|}{u}(x, t)$ .

## *Non-linear heat flow and harmonic maps*

- For functions  $f: M \rightarrow N$  between Riemannian manifolds

$$\Delta f = \text{trace} \nabla df \in \Gamma(f^* TN),$$

i.e.  $(\Delta f)(x) \in T_{f(x)}N$ .

- Heat flow for harmonic maps

$$\begin{cases} \frac{\partial}{\partial t} u = \frac{1}{2} \Delta u \\ u|_{t=0} = u_0 \end{cases}$$

- How to estimate  $|du(t, \cdot)|$ ?

- If  $\text{Sect}^N \leq 0$ , everything works well (Eells-Sampson).
- In general, the heat flow for harmonic maps develops singularities (*blow up in finite time*),  
i.e.  $\exists T > 0, x_0 \in M$ :

$$\limsup_{t \nearrow T} |d u(t, \cdot)|^2(x_0) = \infty$$

- Try to understand how sectional curvature of the target  $N$  enters the estimates of  $|d u(t, \cdot)|$ .



## Damped parallel transport

- **Notation** Let  $X$  be a general semimartingale taking values in a Riemannian manifold  $M$ .

Define the *damped parallel transport*

$$\Theta_t: T_{X_0}M \rightarrow T_{X_t}M$$

by the following covariant equation:

$$\begin{cases} d //_t^{-1} \Theta_t = -\frac{1}{2} //_t^{-1} R^M(\Theta_t, dX_t) dX_t \\ \Theta_0 = \text{id}_{T_{X_0}M} \end{cases}$$

where  $//_t$  is the usual parallel transport along  $X$  with respect to the Levi-Civita connection.

- **Example** Let  $X$  be BM on  $(M, g)$ . Then

$$d //_t^{-1} \Theta_t = -\frac{1}{2} //_t^{-1} \text{Ric}^M(\Theta_t) dt.$$

### Basic observation

- Consider the damped transports

$$\Theta_t^M: T_x M \rightarrow T_{X_t} M, \quad \text{resp.} \quad \Theta_t^N: T_{Y_0} N \rightarrow T_{Y_t} N,$$

along  $X$  on  $M$ , resp. along  $Y$  on  $N$ , where

- $X$  is BM on  $M$  with  $X_0 = x$ , and
  - $Y_t = u(T - t, X_t)$  is the image process on  $N$  for a solution  $u$  of the non-linear heat equation.
- Then

$$\left\{ (\Theta_t^N)^{-1} du(T - t, \cdot)_{X_t(x)} \Theta_t^M : t < T \right\}$$

is a local martingale in  $T_x^* M \otimes T_{u(T,x)} N$ .

## Integration by parts

- In particular, for each  $v \in T_x M$ ,

$$m_t v := (\Theta_t^N)^{-1} du(T - t, \cdot)_{X_t(x)} \Theta_t^M v$$

is a local martingale in  $T_{u(T,x)} N$ .

- Now allow  $v$  to vary with time. For this purpose let  $\ell(t)$  be any adapted process in  $T_x M$  with absolutely continuous paths. Then

$$m_t \ell(t) - \int_0^t m_r d\ell(r)$$

is a local martingale, i.e.,

$$\begin{aligned} & (\Theta_t^N)^{-1} du(T - t, \cdot)_{X_t(x)} \Theta_t^M \ell(t) \\ & - \int_0^t (\Theta_r^N)^{-1} du(T - r, \cdot)_{X_r(x)} \Theta_r^M \dot{\ell}(r) dr \end{aligned}$$

is a local martingale.

## Deformed anti-development

- Consider the process  $\tilde{Y}$ , taking values in  $T_{u(T,x)}N$ , defined by

$$d\tilde{Y}_t = (\Theta_t^N)^{-1} dY_t, \quad \text{where } Y_t = u(T-t, X_t).$$

- Then

$$\tilde{Y}_t = \int_0^t (\Theta_r^N)^{-1} du(T-r, \cdot)_{X_r(x)} //_r^M dB_r$$

where  $B$  is the anti-development of the BM  $X$  in  $T_xM$ , and we see that

$$n_t := (\Theta_t^N)^{-1} du(T-t, \cdot)_{X_t(x)} \Theta_t^M \ell(t) - \tilde{Y}_t \int_0^t \langle \Theta_r^M \dot{\ell}(r), //_r^M dB_r \rangle$$

is a local martingale as well.

- The idea is now to choose  $l_t = \ell(t)$  such that first  $n_t$  is a true martingale and such that

$$l_0 = v, \quad l_\tau = 0 \quad \text{and} \quad \left( \int_0^\tau |\dot{\ell}_t|^2 dt \right)^{1/2} \in L^{1+\varepsilon}$$

(some  $\varepsilon > 0$ ) where, for instance,  $\tau = \tau(x) \wedge T$  with  $\tau(x)$  the first exit time from some rel. compact neighbourhood  $D$  of  $x$ .

- Taking expectations we get

$$\mathbb{E}[n_0] = \mathbb{E}[n_\tau]$$

which gives a gradient formula.

## Theorem (Non-linear gradient formula)

Let  $u : M \times [0, T] \rightarrow N$  be a solution of the non-linear heat equation. Then, for each  $v \in T_x M$ ,

$$du(T, \cdot)_x v = -\mathbb{E} \left[ \tilde{Y}_\tau \int_0^\tau \langle \Theta_s^M \dot{l}_s, \parallel_s dB_s \rangle \right]$$

where:

- $Y_t = u(T - t, X_t)$  and  $\tilde{Y}_t$  its deformed anti-development
- $\tau = \tau(x) \wedge T$  with  $\tau(x) = \inf\{t > 0 : X_t(x) \notin D\}$
- $l_t$  an adapted process with values in  $T_x M$  such that

$$\left( \int_0^\tau |\dot{l}_t|^2 dt \right)^{1/2} \in L^{1+\varepsilon} \quad \text{and} \quad l_0 = v, \quad l_\tau = 0.$$

## Theorem (derivative estimate)

$$|du(T, \cdot)_{xv}| \leq \underbrace{\left\| \int_0^T \langle \Theta_s^M \dot{\ell}_s, //_s^M dB_s \rangle \right\|_p}_{=: \text{(I)}} \cdot \underbrace{\left\| \tilde{Y}_\tau \right\|_q}_{=: \text{(II)}}$$

where  $1 \leq p < \infty$  and  $1/p + 1/q = 1$ .

- **To estimate (I):** Let  $p = 2$  and  $|v| \leq 1$ . Then

$$\text{(I)} \leq c(n) \left( k + \frac{1}{\sqrt{T}} + \frac{1}{r(x)} \right), \quad r(x) := \text{dist}(x, \partial D),$$

if  $\text{Ric}|_D \geq -(n-1)k^2$  with  $k \geq 0$

F.-Y. Wang & A. Th., *JFA* (1998)

- **To estimate (II) :**

Let  $-\kappa_1 \leq \text{Sect}^N \leq \kappa_2$  with  $\kappa_1, \kappa_2 \geq 0$ .

Then

$$\exp\left(-\frac{\kappa_1}{2} \int_0^t h(dY, dY)\right) \leq |(\Theta_t^N)^{-1}| \leq \exp\left(\frac{\kappa_2}{2} \int_0^t h(dY, dY)\right)$$

where  $h$  denotes the metric on  $N$  and  $Y_t := u(T - t, X_t)$ .

M. Arnaudon, X.-M. Li, A. Th., *Ann. Inst. H. Poincaré* (1999)



## Applications *Sharp a priori estimates for harmonic maps*

- Let  $D \subsetneq M$  be a relatively compact domain, and let  $u: D \rightarrow N$  be a harmonic map. Then, for each  $v \in T_x M$ ,  $x \in D$ ,

$$(du)_{xv} = -\mathbb{E} \left[ \widetilde{u(X)}_{\tau} \int_0^{\tau} \langle \Theta_s^M \dot{\ell}_s, //_s dB_s \rangle \right]$$

where  $\tau = \inf\{t > 0 : X_t(x) \notin D\}$ . In particular,

$$|(du)_{xv}| \leq \left\| \int_0^{\tau} \langle \Theta_s^M \dot{\ell}_s, //_s^M dB_s \rangle \right\|_p \cdot \left\| \widetilde{u(X)}_{\tau} \right\|_q$$

- Liouville theorems for harmonic maps*

- Now back to the linear case. The following theorem gives typical gradient formulas for solutions of the heat equation.

### Theorem (Gradient formulas in the linear case)

- Let  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  such that  $\begin{cases} u|_{t=0} = u_0 \\ u(t, \cdot)|_{\partial D} = u_0|_{\partial D}. \end{cases}$  Then

$$du(T, \cdot)_x v = -\mathbb{E} \left[ u_0(X_\tau(x)) \int_0^\tau \langle Q_s \dot{\ell}_s, dB_s \rangle \right], \quad \tau = \tau(x) \wedge T.$$

- Let  $\frac{\partial}{\partial t} u = \frac{1}{2} \Delta u$  such that  $\begin{cases} u|_{t=0} = u_0 \\ u(t, \cdot)|_{\partial D} = 0. \end{cases}$  Then

$$du(T, \cdot)_x v = -\mathbb{E} \left[ u_0(X_T(x)) 1_{\{T < \tau(x)\}} \int_0^{\tau(x) \wedge T} \langle Q_s \dot{\ell}_s, dB_s \rangle \right].$$

## Theorem (Gradient formulas for harmonic functions)

- Let  $\Delta u = 0$  on  $D$ . Then

$$(du)_x v = -\mathbb{E} \left[ u(X_\tau(x)) \int_0^\tau \langle Q_s \dot{\ell}_s, dB_s \rangle \right], \quad \tau = \tau(x).$$

- If  $\text{Ric}|_D \geq -(n-1)k^2$  for some  $k \geq 0$ , it is easy to derive from here:

$$|\nabla u|(x) \leq \|u\|_D \left\| \int_0^\tau \langle Q_r \dot{\ell}_r, \ell_r dB_r \rangle \right\|_2 \leq \|u\|_D c(n) \left( k + \frac{1}{r(x)} \right)$$

where  $r(x) := \text{dist}(x, \partial D)$ .

- But recall *Cheng-Yau*! For  $u$  **positive** we want:

$$|\nabla u|(x) \leq u(x) \cdot c(n) \left( k + \frac{1}{r(x)} \right).$$

## Problem

$$\frac{|\nabla u|}{u}(x) = -\mathbb{E} \left[ \underbrace{\frac{u(X_\tau(x))}{u(x)}}_{L^1 ?} \underbrace{\int_0^\tau \langle Q_s \dot{\ell}_s, dB_s \rangle}_{L^\infty ?} \right]$$

Note:  $Y_t := \frac{u(X_t(x))}{u(x)}$  is a strictly positive martingale;  $Y_0 = 1$ .

## Lemma

Let  $Y$  be a strictly positive martingale;  $Y_0 = y$

$$\implies \forall \alpha \in ]0, 1[ : \quad \mathbb{E}[\langle Y, Y \rangle_\infty^{\alpha/2}]^{1/\alpha} \leq C_\alpha y.$$

- Proof. Without restriction  $Y_0 = 1$

$$Y = 1 + \beta_{\langle Y, Y \rangle}, \quad \beta \text{ real BM (with } \beta_0 = 0)$$

$$\implies \langle Y, Y \rangle \leq S := \inf\{s > 0 : \beta_s = -1\}$$

$$\text{But } S \sim 1/\beta_1^2$$

$$\implies \mathbb{E}[S^{\alpha/2}]^{1/\alpha} = \mathbb{E}[|\beta_1|^{-\alpha}]^{1/\alpha} =: C_\alpha = \frac{1}{\sqrt{2}} \left( \frac{\Gamma(\frac{1-\alpha}{2})}{\Gamma(\frac{1}{2})} \right)^{1/\alpha}.$$

- Here,

$$\forall \alpha < 1 : \quad \mathbb{E} \left[ \left( \int_0^\tau |du|^2(X_r) dr \right)^{\alpha/2} \right]^{1/\alpha} \leq C_\alpha u(x)$$

$$\text{et } \mathbb{E}[u(X_\tau)^\alpha]^{1/\alpha} \leq C_\alpha u(x).$$

- Let  $n_r := \frac{\nabla u}{|\nabla u|}(X_r)$

Theorem (M. Arnaudon, B. Driver, A. Th., 2006)

For each  $\alpha \geq \frac{n-2}{n-1}$ ,

$$N_s := |du|^\alpha(X_s) \exp\left(-\frac{\alpha}{2} \int_0^r \text{Ric}(n_r, n_r) dr\right)$$

is a (local) submartingale.

- **Consequence**  $N_s l_s - \int_0^s N_r dl_r$  submartingale if  $\dot{l}_s \leq 0$   
 [Here:  $l_s$  scalar,  $l_0 = 1$ ,  $l_\tau = 0$  with  $\tau = \tau(x)$ ]

### Corollary (Inequalities of Bismut type)

$$|du|^\alpha(x) \leq -\mathbb{E} \left[ \int_0^T N_r \dot{\ell}_r dr \right], \quad \alpha \geq \frac{n-2}{n-1}.$$

By Hölder, if  $\alpha < 2$ ,

$$\begin{aligned} |du|^\alpha(x) &\leq \mathbb{E} \left[ \left( \int_0^T |du|^2(X_s) ds \right)^{\alpha/2} \right. \\ &\quad \left. \times \left( \int_0^T \exp \left\{ \frac{\alpha}{\alpha-2} \int_0^s \text{Ric}(n_r, n_r) dr \right\} |\dot{\ell}_s|^{2-\alpha} ds \right)^{\frac{2-\alpha}{2}} \right]. \end{aligned}$$

Let  $\alpha \in \left[\frac{n-2}{n-1}, 1\right[$  if  $n \geq 3$ , and  $\alpha \in ]0, 1[$  if  $n = 2$ ,  
 $p > 1$  such that  $\alpha p < 1$ ,  
 $q > 1$  conjugated exponent to  $p$ .

Suppose that  $\text{Ric} \geq -K$  for some  $K \geq 0$ . Then,

$$|\nabla u|(x) \leq C_{\alpha p} u(x) \mathbb{E} \left[ \left( \int_0^T \exp \left\{ \frac{\alpha}{2-\alpha} Ks \right\} |\dot{\ell}_s|^{\frac{2}{2-\alpha}} ds \right)^{\frac{(2-\alpha)q}{2}} \right]^{\frac{1}{\alpha q}}.$$

### Corollary

$$|\nabla \log u|(x) \leq c(n) \left[ k + \frac{1}{r(x)} \right]$$

if  $\text{Ric}^M \geq -(n-1)k^2$ ,  $k \geq 0$ .



## From gradient estimates to Harnack inequalities

An obvious consequence of the Cheng-Yau estimate is:

### Corollary (Harnack inequality)

Let  $u$  be harmonic on  $B_r(x) \subset M$  where  $M$  is complete. Then

$$\sup_{B_{r/2}(x)} u \leq c(n, r, k) \inf_{B_{r/2}(x)} u$$

- **Elliptic case** Let  $u: D \rightarrow \mathbb{R}$  be harmonic and  $x_1, x_2 \in D$ . Fix  $B \subset D$  open, rel. compact, connected, such that  $x_1, x_2 \in B$ .

Then  $\frac{u(x_1)}{u(x_2)} \leq C$  where the constant  $C$  depends only on

- the lower bound for Ric on  $B$ ;
- $\text{dist}_B(x_1, x_2)$  and  $\text{dist}(x_i, \partial B)$ ,  $i = 1, 2$ ;
- $n = \dim M$ .
- **Parabolic case** Want to compare  $P_T f(x_1)$  and  $P_T f(x_2)$

- **Harnack inequality with power**  $\alpha > 1$  (F.-Y. Wang)

$$(P_T f)^\alpha(x) \leq (P_T f^\alpha)(y) \cdot C(T, K, d(x, y), \alpha)$$

- *Idea* Estimate the derivative of

$$[0, 1] \ni r \mapsto \log(P_T f^{\beta(r)})^{\alpha(r)}(\gamma(r))$$

$$\text{where } \gamma(0) = x, \gamma(1) = y$$

$$\alpha(0) = \alpha, \alpha(1) = 1$$

$$\beta(0) = 1, \beta(1) = \alpha$$

- If  $B$  is an open relatively compact connected neighbourhood of  $x, y$ , the constant should depend only on a lower bound of  $Ric$  on  $B$ .

- Need local gradient estimates of the type:

$$\frac{|\nabla P_T f(x)|}{P_T f(x)} \leq \frac{C_1}{T \wedge 1} + C_2 P_T \left( \frac{f}{P_T f(x)} \log \frac{f}{P_T f(x)} \right)$$

- For  $u = P_T f > 0$  use

- $(\frac{1}{2}\Delta - \partial_t) \frac{|\nabla u|^2}{u} \geq \frac{1}{u} \text{Ric}(\nabla u, \nabla u) \geq -K \frac{|\nabla u|^2}{u}$

- $(\frac{1}{2}\Delta - \partial_t) u \log u = \frac{1}{2} \frac{|\nabla u|^2}{u}$

- Then, if  $\text{Ric} \geq -K$ ,

$$N_t := \frac{1}{2} \frac{T-t}{1+K(T-t)} \frac{|\nabla P_{T-t} f|^2}{P_{T-t} f}(X_t) + (P_{T-t} f \log P_{T-t} f)(X_t)$$

is a local submartingale and  $\mathbb{E}[N_0] \leq \mathbb{E}[N_T]$  gives an inequality of the wanted type ( $\rightarrow$  localization).

### Corollary (Wang-Arnaudon-A.Th. 2007)

*Let  $M$  be an arbitrary complete Riemannian manifold.*

*Then, for any  $\delta > 2$  there exists a positive function*

*$C_\delta \in C([0, \infty[ \times M)$  such that the transition density  $p_t(x, y)$  of  $P_t$  with respect to the volume measure satisfies*

$$p_t(x, y) \leq \frac{\exp \left\{ -\text{dist}(x, y)^2 / (2\delta t) + C_\delta(t, x) + C_\delta(t, y) \right\}}{\sqrt{\text{vol}(B(x, \sqrt{2t}))\text{vol}(B(y, \sqrt{2t}))}},$$

$x, y \in M, t \in ]0, \infty[.$